Network Systems
Science & Advanced
Computing

Biocomplexity Institute & Initiative

University of Virginia

Estimation of COVID-19 Impact in Virginia

May 18th, 2022

(data current to May 15th – May 17th)
Biocomplexity Institute Technical report: BI-2022-1439



BIOCOMPLEXITY INSTITUTE

biocomplexity.virginia.edu

About Us

- Biocomplexity Institute at the University of Virginia
 - Using big data and simulations to understand massively interactive systems and solve societal problems
- Over 20 years of crafting and analyzing infectious disease models
 - Pandemic response for Influenza, Ebola, Zika, and others



Points of Contact

Bryan Lewis brylew@virginia.edu

Srini Venkatramanan srini@virginia.edu

Madhav Marathe marathe@virginia.edu

Chris Barrett@virginia.edu

Model Development, Outbreak Analytics, and Delivery Team

Przemyslaw Porebski, Joseph Outten, Brian Klahn, Alex Telionis, Srinivasan Venkatramanan, Bryan Lewis, Aniruddha Adiga, Hannah Baek, Chris Barrett, Jiangzhuo Chen, Patrick Corbett, Stephen Eubank, Galen Harrison, Ben Hurt, Dustin Machi, Achla Marathe,

Madhav Marathe, Mark Orr, Akhil Peddireddy, Erin Raymond, James Schlitt, Anil Vullikanti,

Lijing Wang, James Walke, Andrew Warren, Amanda Wilson, Dawen Xie



Overview

• Goal: Understand impact of COVID-19 mitigations in Virginia

Approach:

- Calibrate explanatory mechanistic model to observed cases
- Project based on scenarios for next 4 months
- Consider a range of possible mitigation effects in "what-if" scenarios

Outcomes:

- Ill, Confirmed, Hospitalized, ICU, Ventilated, Death
- Geographic spread over time, case counts, healthcare burdens

Key Takeaways

Projecting future cases precisely is impossible and unnecessary. Even without perfect projections, we can confidently draw conclusions:

- Case rates and hospitalizations continue to plateau
- VA 7-day mean daily case rate continues to increase to 32/100K from 26/100K
 - US also increased to 36/100K (from 22/100K)
- District level fitting seems to have helped with the projection's accuracy
- BA.2.12.1 continues to rise, but pace of growth remains slow and steady


The situation continues to change. Models continue to be updated regularly.

20-May-22

Situation Assessment



Case Rates (per 100k) and Test Positivity




District Trajectories

Goal: Define epochs of a Health District's COVID-19 incidence to characterize the current trajectory

Method: Find recent peak and use hockey stick fit to find inflection point afterwards, then use this period's slope to define the trajectory

Hockey stick fit




Trajectory	Description	Weekly Case Rate (per 100K) bounds
Declining	Sustained decreases following a recent peak	below -0.9
Plateau	Steady level with minimal trend up or down	above -0.9 and below 0.5
Slow Growth	Sustained growth not rapid enough to be considered a Surge	above 0.5 and below 2.5
In Surge	Currently experiencing sustained rapid and significant growth	2.5 or greater




District Trajectories – last 10 weeks

Status	# Districts (prev week)
Declining	0 (2)
Plateau	4 (0)
Slow Growth	7 (11)
In Surge	24 (22)

Curve shows smoothed case rate (per 100K) Trajectories of states in label & chart box Case Rate curve colored by Reproductive number





MUNIVERSITY of VIRGINIA


■ 1.5 <= R < 2

CDC's new COVID-19 Community Levels

What Prevention Steps Should You Take Based on Your COVID-19 Community Level?

Low	Medium	High		
 Stay <u>up to date</u> with COVID-19 vaccines <u>Get tested</u> if you have symptoms 	 If you are at high risk for severe illness, talk to your healthcare provider about whether you need to wear a mask and take other precautions Stay up to date with COVID-19 vaccines Get tested if you have symptoms 	 Wear a mask indoors in public Stay up to date with COVID-19 vaccines Get tested if you have symptoms Additional precautions may be needed for people at high risk for severe illness 		
People may choose to mask at any time. People with symptoms, a positive test, or exposure to someone with COVID-19				

COVID-19 Community Levels – Use the Highest Level that Applies to Your Community					
New COVID-19 Cases Per 100,000 people in the past 7 days	Indicators	Low	Medium	High	
Fewer than 200	New COVID-19 admissions per 100,000 population (7-day total)	<10.0	10.0-19.9	≥20.0	
	Percent of staffed inpatient beds occupied by COVID-19 patients (7-day average)	<10.0%	10.0-14.9%	≥15.0%	
200 or more	New COVID-19 admissions per 100,000 population (7-day total)	NA	<10.0	≥10.0	
	Percent of staffed inpatient beds occupied by COVID-19 patients (7-day average)	NA	<10.0%	≥10.0%	




The COVID-19 community level is determined by the higher of the new admissions and inpatient beds metrics, based on the current level of new cases per 100,000 population in the past 7 days



CDC Data Tracker Portal

should wear a mask.

CDC's new COVID-19 Community Levels



Red outline indicates county had 200 or more cases per 100k in last week


Pale color indicates either beds or occupancy set the level for this county

Dark color indicates both beds and occupancy set the level for this county

COVID-19 Community Levels – Use the Highest Level that Applies to Your Community					
New COVID-19 Cases Per 100,000 people in the past 7 days	Indicators	Low	Medium	High	
Fewer than 200	New COVID-19 admissions per 100,000 population (7-day total)	<10.0	10.0-19.9	≥20.0	
	Percent of staffed inpatient beds occupied by COVID-19 patients (7-day average)	<10.0%	10.0-14.9%	≥15.0%	
200 or more	New COVID-19 admissions per 100,000 population (7-day total)	NA	<10.0	≥10.0	
	Percent of staffed inpatient beds occupied by COVID-19 patients (7-day average)	NA	<10.0%	≥10.0%	

The COVID-19 community level is determined by the higher of the new admissions and inpatient beds metrics, based on the current level of new cases per 100,000 population in the past 7 days

District Trajectories with Community Levels





Estimating Daily Reproductive Number –

Redistributed gap

May 13th Estimates

Region	Date Confirmed R _e	Date Confirmed Diff Last Week
State-wide	1.111	0.030
Central	1.096	-0.020
Eastern	1.090	0.065
Far SW	1.026	0.017
Near SW	1.124	0.084
Northern	1.115	0.025
Northwest	1.155	0.049






Skipping Weekend Reports & holidays biases estimates
Redistributed "big" report day to fill in gaps, and then estimate R from
"smoothed" time series

Methodology

- Wallinga-Teunis method (EpiEstim¹) for cases by confirmation date
- Serial interval: Discrete distribution from observations (mean=4.3, Flaxman et al, Nature 2020)
- Using Confirmation date since due to increasingly unstable estimates from onset date due to backfill


1. Anne Cori, Neil M. Ferguson, Christophe Fraser, Simon Cauchemez. A New Framework and Software to Estimate Time-Varying Reproduction Numbers During Epidemics. American Journal of Epidemiology, Volume 178, Issue 9, 1 November 2013, Pages 1505–1512, https://doi.org/10.1093/aje/kwt133



Mask Usage and Vaccination

Self-reported mask usage continues to fall

- US and VA continue to slightly decrease
- Mask wearing remains lower amongst unvaccinated, especially among least willing to be vaccinated






Data Source: https://covidcast.cmu.edu

Wastewater Monitoring

Wastewater provides a coarse early warning of COVID-19 levels in communities


- Overall in the US, there is an increase in sites with increased levels of virus compared to 15 days ago
- Proportion of sites with current virus levels at or exceeding the max of previous historical levels, has further increased since last week




COVID-like Illness Activity

COVID-like Illness (CLI) gives a measure of COVID transmission in the community

- Emergency Dept (ED) based CLI is more correlated with case reporting
- Urgent Care (UC) is more sensitive and is a leading indicator but is prone to some false positives
- As testing behaviors and case ascertainment levels shift, these measures may capture disease better than confirmed cases
- Current trends in UC CLI are slightly up





SARS-CoV2 Variants of Concern

Emerging new variants will alter the future trajectories of pandemic and have implications for future control


- Emerging variants can:
 - Increase transmissibility
 - Increase severity (more hospitalizations and/or deaths)
 - Limit immunity provided by prior infection and vaccinations
- Genomic surveillance remains very limited
 - Challenges ability to estimate impact in US to date and estimation of arrival and potential impact in future

WHO label	Pango lineage•	GISAID clade	Nextstrain clade	Additional amino acid changes monitored°	Earliest documented samples	Date of designation
Alpha	B.1.1.7	GRY	20I (V1)	+S:484K +S:452R	United Kingdom, Sep-2020	18-Dec-2020
Beta	B.1.351	GH/501Y.V2	20H (V2)	+S:L18F	South Africa, May-2020	18-Dec-2020
Gamma	P.1	GR/501Y.V3	20J (V3)	+S:681H	Brazil, Nov-2020	11-Jan-2021
Delta	B.1.617.2	G/478K.V1	21A, 21I, 21J	+S:417N +S:484K	India, Oct-2020	VOI: 4-Apr-2021 VOC: 11-May-2021
Omicron*	B.1.1.529	GRA	21K, 21L	+R346K	Multiple countries, Nov-2021	VUM: 24-Nov-2021 VOC: 26-Nov-2021


Omicron Prevalence

CDC nowcast for week ending April 2nd shows 98% overall BA.2 in Region 3 with BA 2.12.1 at 56% (last week 48%)

Overall BA.2 in USA now at 99% (BA.2.12.1 at 37%, last week 27%)



HHS Region 3: 2/6/2022 - 5/14/2022





HHS Region 3: 5/8/2022 - 5/14/2022 NOWCAST

Enumerated lineages are US VOC and lineages circulating above 1% nationally in at least one week period. *Other* represents the aggregation of lineages which are circulating 1% nationally during all weeks displayed.
 These data include Nowcast estimates, which are modeled projections that may differ from weighted estimates operated at later dates.


A11-141-133 and trier subinsages are aggregated with 5.1617.2 DA1, 1 BA3, BA4, BA5 and their subinsages (except BA1.1 and its subinsages) are aggregated with B1.1.529, For regional data, BA.1.1 and its subinsages are also aggregated with B1.1.529, as they currently cannot be reliably called in each region. Except BA2.12.1 and its sublineages, BA2 sublineages are aggregated with BA2.

Collection date, week ending





SARS-CoV2 Omicron and Sub-Variants

As detected in whole Genomes in public repositories



VoC Polynomial Fit Projections







Note: Data lags force projections to start in past. Everything from dotted line forward is a projection.

GISAID

Pandemic Pubs

- **1.** New data gives estimates on the degree of protection from post-acute sequelae PASC offered by vaccines.
- **2.** European study gives baseline risk of 1% of venous or arterial thromboembolism among COVID patients. Increases among males, hospitalizations and the elderly.
- **3.** ECDC list BA.4 and BA.5 as VoCs. Cases appear to have peaked in S. Africa with low severe outcomes.
- **4.** UK: Bank of England's Monetary Policy Committee reports concern that share of the 16-64 y/o population who are outside the workforce and do not seeking a job because of long-term sickness is a record high.






May 2021 Sep 2021

Sep 2020 Patients with confirmed COVID-19 diagnosis, minimum age of 18 years with 3 month follow-up post-diagnosis between September 21, 2020 and December 14, 2021 were identified from TriNetX research network platform. 1,578,719 patients with confirmed COVID-19 were identified and 1.6% (n=25,225) completed vaccination. Before matching vaccinated individuals who were found to be positive tended to be older and have more

comorbidities. https://academic.oup.com/ofid/advancearticle/doi/10.1093/ofid/ofac228/6582238?login=false#.Yn2vXtn7Cpg.t witter



Tulio de Oliveira Director of CERI (Centre for Epidemic Response & Innovatior S.Africa) reports BA.4 and BA.5 have significantly increased cases in countries with strong BA.1 but minimal BA.2 waves, S. Africa and Portugal. Locations w similar conditions could be more vulnerable to these immune escape variants https://twitter.com/Tuliodna/status/1526470866762141697







Table 2. Post-Acute Sequelae of Sars-Cov-2 (PASC) Mortality and Morbidity Risk at 28 days: vaccine vs. no-vaccine.





primary care records from the Netherlands, Italy, Spain, and the UK, and outpatient specialist records from Germany. Overall, 909 473 COVID-19 cases and 32329 patients hospitalised with COVID-19 on or after Sept 1. 2020, were studied. Cumulative 90-day incidence of venous thromboembolism ranged from 0.2% to 0.8% among COVID-19 cases, and up to 4.5% for those hospitalised. The occurrence of venous thromboembolism in patients with COVID-19 was associated with an increased risk of death (adjusted HRs 4-42 [3.07-6.36] for those not hospitalised and 1.63 [1·39-1·90] for those hospitalised), as was the occurrence of arterial thromboembolism (3.16 [2.65-3.75] and 1.93 [1.57-2.37]). https://www.thelancet.com/journals/laninf/art icle/PIIS1473-3099(22)00223-7/fulltext


Epidemic and Variant Dynamics in South Africa



United States Overall


- Nation pivoting towards more growth, focused in Northeast
- Most are sustained declines

Status	# States
Declining	7 (8)
Plateau	18 (22)
Slow Growth	7 (6)
In Surge	22 (18)



Virginia and Her Neighbors

- More growth experienced across neighbors
- Some states at very high levels (CT, NJ)
- Neighbors to south starting to ramp up as well




United States Hospitalizations

 Hospital admissions are lagging case rates


Status	# States (prev week)		
Declining	1 (1)		
Plateau	17 (21)		
Slow Growth	28 (23)		
In Surge	7 (5)		


Trajectories of States



County-level comparison to previous highest peak

- Most counties in VA have had the highest case rate of the pandemic in the last week
- Nationally the number of counties at their highest rate has expanded considerably







County-level comparison to last Summer

- Most counties in VA have had the highest case rate of the pandemic in the last week
- Nationally the number of counties at their highest rate has expanded considerably

Recent Incidence Compared to Weekly Summer Mean by County Mean: 14.85; Median: 0.87; IQR: 0.25-2.64







Using Ensemble Model to Guide Projections

Ensemble methodology that combines the Adaptive with machine learning and statistical models such as:


- Autoregressive (AR, ARIMA)
- Neural networks (LSTM)
- Kalman filtering (EnKF)


Weekly forecasts done at county level.


Models chosen because of their track record in disease forecasting and to increase diversity and robustness.


Ensemble forecast provides additional 'surveillance' for making scenario-based projections.


Also submitted to CDC Forecast Hub.












Last week's projection comparison


Projection from last week (May 7th)

Virginia Daily Confirmed - Comparison 2022-05-07



Projection from 3 weeks ago (April 23rd)

Virginia Daily Confirmed - Comparison 2022-04-23



Additional Analyses





COVID-19 Scenario Modeling Hub – Round 13

Collaboration of multiple academic teams to provide national and state-by-state level projections for 4 aligned scenarios

- Round 13 results getting finalized
 - Scenarios: New Variant in Summer and waning compared (yes/no new variant vs. 4 month or 10 month waning)
- Prelim results shared internally
- Only national consortium tracking Omicron wave well
- Rounds 4-12 now available Round 4 Results were published
 May 5th, 2021 in MMWR

https://covid19scenariomodelinghub.org/viz.html





20-May-22 27